

Quantum Hall effect in honeycomb lattice under strain

Abolhassan Vaezi,^{1,2,*} Nima Abedpour,¹ and Reza Asgari^{1,†}

¹*School of Physics, Institute for Research in Fundamental Sciences, IPM, Tehran, 19395-5531, Iran*

²*Department of Physics, Sharif University of Technology, Tehran 11155-9161, Iran*

We investigate the possibility of the quantum Hall effect in the deformed honeycomb lattice. Static strain in the honeycomb lattice preserves the time reversal symmetry (TRS), and can be viewed as a *valley dependent* gauge field in the continuum limit. We calculate the response of the system to the *valley magnetic field* induced by strain and compute the *valley Hall conductance*. It is shown that the system exhibits a nontrivial response (nonzero Hall conductance) even in time reversal invariant (TRI) samples. We argue that a quantized current will be induced if we subject the system to a *valley electric field* by applying time dependent strain. A theory for the low energy excitations of the systems is developed. We show that the mutual Chern-Simons action describes the low energy theory of this system and also discuss the necessary details for observing such a behavior with a proposal for experiment.

PACS numbers: 73.43.Cd, 61.48.Gh, 81.40.Jj, 07.55.Db

Introduction.— The emergence of the fractional charge and statistics in quantum Hall systems, has spurred a broad interest in topological states of matter. In semiconductor quantum Hall liquids, strong magnetic field at integer and fractional filling fractions can generate quantized off-diagonal conductivity [1]. The experimental discovery of the quantum Hall effect (QHE) in monolayer graphene, has attracted a lot of attention to the properties of two-dimensional (2D) Dirac fermions in the presence of magnetic field [2]. Haldane in a seminal paper [3] showed that in order to have integer quantum Hall effect on honeycomb lattice, the presence of the external magnetic field is not necessary. He suggested a model in which the time reversal symmetry (TRS) breaks because of the pattern of the hopping integrals, and the Hall conductance does not vanish despite the vanishing total magnetic field. Accordingly, it is believed that the time-reversal symmetry breaking by itself can potentially lead to integer QHE. This has been recently verified for partial filling fractions as well [4]. It is also well established that topological insulators which preserve TRS exhibit quantum *spin* Hall effect [5, 6]. In this paper we discuss that static strain which does not break time reversal symmetry induces a nonzero *valley magnetic field* whose direction changes as we travel from one valley to another in the honeycomb lattice. We show that this valley magnetic field induces carrier density with a quantized valley Hall conductance. We discuss through the Laughlin argument and the Chern-Simons theory that a time dependent strain which induces a nonzero *valley electric field* as well, generates a nonzero current in the system. It is also shown that the low-energy behavior of this system can be described by the mutual $U(1) \times U(1)$ Chern-Simons action [7].

In the following, we discuss a possible realization for suggestion for charge quantum Hall effect (CQHE). In the honeycomb lattice with massive Dirac fermions, for each spin degree of freedom, the low energy description is




FIG. 1: (Color online) (a): Upper hemisphere represents the Berry curvature for $H_{K,s}$ which leads to $C_K = C_{K,\uparrow} + C_{K,\downarrow} = 1$. Similarly, lower hemisphere represents $H_{K',s}$. According to Eq. (1), the pseudo-magnetic field induces the same charge density at K and K' valleys. (b): The sign of the spatial current along the x -axes is determined by $C\tilde{E}_y$. Hence, K and K' contribute the same current.

given by the sum of two independent sub-Hamiltonians. We can tune the structure of these sub-Hamiltonians to have opposite Chern numbers. On the other hand, it has been discussed in the literature that strain on honeycomb lattice can induce an effect similar to that of the magnetic field, which is called *pseudo-magnetic* field [8]. Basically, strains changes the bond length between atoms and due to the particular band structure and lattice symmetry of the honeycomb lattice, strain creates pseudo-magnetic fields acting on the charge carriers. Because strain does not break TRS, the two sub-Hamiltonians feel opposite pseudo-magnetic fields. Massive pseudo-magnetic fields raised by highly strained nano-bubbles that form when graphene is grown on a platinum sur-

face have been recently measured by STM spectroscopic measurements from the spectral peak spacing [9]. The spectral peak indicates Landau levels created by the local pseudo-magnetic fields. Accordingly, strain engineering can prepare the desired system.

Since spatial current of edge states in the QHE is determined by the product of the Chern number and the sign of the pseudo-electric field [10], both bands contribute the same current and therefore the total current is nonzero. More specifically, let H_1 and H_2 be Hamiltonians with Chern numbers C_1 and C_2 , respectively. We assume that H_1 and H_2 are subject to $\tilde{\mathbf{B}}_1$ and $\tilde{\mathbf{B}}_2$ pseudo-magnetic fields. Then the conductivity is determined by $J_{i,x} = \frac{e^2}{h} C_i \tilde{E}_{i,y}$ where $i \in \{1, 2\}$, and $J_{i,x}$ is the x -component of the current induced by the pseudo-electric field $\tilde{E}_{i,y}$. The current thus depends on $C_i \tilde{E}_{i,y}$. Requiring the time reversal invariance of $H_1 \oplus H_2$, we obtain $C_2 = -C_1$ and $\tilde{\mathbf{B}}_2 = -\tilde{\mathbf{B}}_1$, whence $\tilde{\mathbf{A}}_2 = -\tilde{\mathbf{A}}_1$. Therefore $\tilde{E}_{2,y} = -\tilde{E}_{1,y}$, and the total current $J_x = J_{1,x} + J_{2,x}$ and the quasiparticle density are given by:

$$J_x = \frac{e^2}{h} (C_1 \tilde{E}_{1,y} + C_2 \tilde{E}_{2,y}), \quad n = \frac{e^2}{h} (C_1 \tilde{B}_{1,z} + C_2 \tilde{B}_{2,z}) \quad (1)$$

It means that those circumstances are sufficient to have a nonzero total edge current. There is also another way to justify the above claim. Within the continuum model, H_1 and H_2 are independent and are invariant under global $U(1)$ gauge transformations. The corresponding Noether currents are $\mathbf{J}_1$ and $\mathbf{J}_2$. Therefore, both $\mathbf{J}_+ = \mathbf{J}_1 + \mathbf{J}_2$ and $\mathbf{J}_- = \mathbf{J}_1 - \mathbf{J}_2$ are conserved quantities [11]. $\mathbf{J}_+$ is the electromagnetic response of the system, while $\mathbf{J}_-$ is the response to the pseudo-electromagnetic field. To this end, note that everywhere $q_i \tilde{\mathbf{A}}_i$ appears in the Hamiltonian where q_i is a particle charge. We can safely assume $\tilde{\mathbf{A}}_2 = \tilde{\mathbf{A}}_1$ and $q_2 = -q_1$. Consequently, $\tilde{\mathbf{A}}$ acts like ordinary electromagnetic gauge field which couples to opposite charges at two valleys. This is a way to detect the conserved current $\mathbf{J}_-$, which can in principle be nonzero.

In Ref. [12], a condition to determine a zero-field QHE in massless graphene and quantum *pseudo-spin* Hall effect like topological insulators in the presence of *pseudo-spin*-orbit coupling have been proposed. Nonetheless, since the two generated edge states here circulate in opposite direction, the net current would be zero; thus instead of a CQHE a pseudo-spin QHE would emerge.

Physical realization.— Let us consider the Hamiltonian of whether undoped graphene or an artificial 2D honeycomb lattices in the continuum approximation [3, 13]. In that limit it is given by the sum of the following two independent massless Dirac Hamiltonians ($\hbar = 1$):

$$H_{Q,s} = v_F \int d^2x \psi_{Q,s}^\dagger(x) (\pm i\sigma^x \partial_x + i\sigma^y \partial_y) \psi_{Q,s}(x) \quad (2)$$

where $Q = \{K, K'\}$ and $\sigma^{x,y,z}$ are Pauli matrices, K and K' refers to two Dirac points and s is the spin degree of freedom. The sign $+$ ($-$) refers to K (K'), respectively. Because we are interested in the Hall conductance of the system, we need to compute the Chern number of the ground state. We require the resulting Hamiltonian i.e. $H_K \oplus H_{K'}$ to be time reversal as well as spin rotation invariant. To this end, we do need an opening gap mechanism in the system. In this case, the mass term for both valleys should be equal and independent of the spin of electrons. Therefore, we should have:

$$H_{Q,\sigma} \rightarrow H_{Q,\sigma} + \int d^2x \psi_{Q,\sigma}^\dagger(x) [m\sigma^z] \psi_{Q,\sigma}(x) \quad (3)$$

It is easy to check that the above Hamiltonian neither breaks the time reversal nor the spin rotation symmetry. A generic noninteracting two band Hamiltonian can be written as $H = -\sum_k \psi^\dagger(k) \left(d_0(k) \mathbf{1} + \vec{d}(k) \cdot \vec{\sigma} \right) \psi(k)$. For every k , the term in parenthesis resembles the Hamiltonian for a spin 1/2 in a magnetic field given by $\vec{d}_k$, and therefore, $\psi(k)$ aligns itself with the direction of $\vec{d}_k$. Thus the Hamiltonian maps the k space (2D torus) to a unit 2D sphere spanned by $\vec{d}_k$. The Chern number can be viewed as the winding number of this mapping [7, 10]. It can also be computed through $C = \frac{\theta_B}{2\pi}$, where θ_B is the Berry phase of the ground state $|\hat{d}_k\rangle$ along the path given by $(0, 0) \rightarrow (2\pi, 0) \rightarrow (2\pi, 2\pi) \rightarrow (0, 2\pi) \rightarrow (0, 0)$. Using this recipe, it can be shown that the Chern number of the massive Dirac Hamiltonian of the following form

$$H = \int d^2x \psi_{K,\sigma}^\dagger (m\sigma^z + i\eta_1 \sigma^x \partial_x + i\eta_2 \sigma^y \partial_y) \psi_{K,\sigma} \quad (4)$$

is given by $C = \frac{\theta_B}{2\pi} = \frac{1}{2} \text{sgn}(\eta_1 \eta_2 m)$ [7]. Therefore, we have:

$$C_K = \sum_\sigma C_{K,\sigma} = +1, \quad C_{K'} = \sum_\sigma C_{K',\sigma} = -1 \quad (5)$$

Because we require the system to be time reversal and spin rotation invariant, the mass term should be independent of the valley degree of freedom as well as the spin of quasiparticles. One possible way to generate such a mass term is to introduce staggered chemical potential i.e. on sublattice A we have $\mu_A = m$ and on sublattice B we have $\mu_B = -m$. This pattern of the chemical potential generates a mass term as we desire. This mechanism, for instance can be realized experimentally in graphene by chemical doping [14, 15].

Another mechanism to generate the mass term is to take strong on-site repulsions, U , into account e.g. by adding the Hubbard term. The system is no longer graphene in that case, but still lives on honeycomb lattice [16]. In the rest of this paper we will demonstrate

that our result depends only on the honeycomb lattice structure and applies to this case as well. Several authors including one of us, have studied the properties of the Hubbard model near the Mott transition. Both numerical and theoretical studies have reported the existence of *gapped* spin liquids [17–20]. In this state of matter, the fermionic degree of freedom can be modeled by two gapped Dirac Hamiltonians for each spin with a mass term of the form we used in the previous section. Therefore, slightly above the Mott transition ($U > U_c$), but still below the onset of anti-ferromagnetic order, a mass term will be added to the system which respects all symmetries of the lattice including time reversal, spin rotation, parity, 60° rotation and translation symmetries [21].

Strain engineering and induced pseudo-magnetic field.— It is interesting to investigate the effect of strain on defect-free honeycomb lattices due to smooth elastic deformations. Strains can create induce gauge field and of at least 10% can be induced in the system without damaging its structure [22]. Importantly, due to the gauge field of strains, quasiparticles near K point feel a pseudo-magnetic field that is opposite to the pseudo-magnetic field that quasiparticles near K' feel, in agreement with the requirement of the time reversal invariance. Therefore we replace H_K and $H_{K'}$ with the following Hamiltonians, respectively ($e = 1$):

$$\begin{aligned} H_{K,\sigma} &= \int d^2x \psi_{K,\sigma}^\dagger \left[m\sigma^z + v_F \sigma^x (i\partial_x - \tilde{A}_x) \right. \\ &\quad \left. + v_F \sigma^y (i\partial_y - \tilde{A}_y) \right] \psi_{K,\sigma} \\ H_{K',\sigma} &= \int d^2x \psi_{K',\sigma}^\dagger \left[m\sigma^z - v_F \sigma^x (i\partial_x + \tilde{A}_x) \right. \\ &\quad \left. + v_F \sigma^y (i\partial_y + \tilde{A}_y) \right] \psi_{K',\sigma} \end{aligned} \quad (6)$$

$\tilde{\mathbf{A}}$ connects to the strain of the sample like [23, 24]

$$\tilde{\mathbf{A}} = \frac{\Phi_0 c \beta}{a} \begin{pmatrix} u_{xx} - u_{yy} \\ -2u_{xy} \end{pmatrix} \quad (7)$$

where u_{ij} is the strain distribution, $\beta \approx \partial \log(t)/\partial \log(a) \approx 2$, $a \approx 1.4 \text{ \AA}$ is the lattice constant, $t \approx 2.7 \text{ eV}$ is the nearest neighbor coupling energy, and $c = \sqrt{3}\kappa/(2\pi)$ is a constant and $\kappa \approx 0.56$ [25]. Φ_0 is the quantum unit of magnetic flux. We will propose a feasible way to generate proper strains and pseudo-magnetic fields on strained graphene nanoribbons.

Now at this stage, we use auxiliary gauge field, $a_{Q,\mu}$ to write the current of quasiparticles living on the Dirac cone at valley Q (where $Q = \{K, K'\}$) as $j_Q^\mu = \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \partial_\nu a_{Q,\rho}$ ($\hbar = 1$, $e = 1$). The reason is that $\partial_\mu j_Q^\mu = 0$ is naturally satisfied using that definition. We can obtain the effective action for the gauge field degree of freedom, which describes the low energy theory of the system under consideration, by integrating out fermions. Because the Chern number for both H_K and $H_{K'}$ is non-zero, it is known that the Chern-Simons action describes

the effective low energy theory for both, and therefore the Lagrangian corresponding to H_Q is:

$$\mathcal{L}_Q = \pm \frac{1}{4\pi} \epsilon^{\mu\nu\rho} a_{Q,\mu} \partial_\nu a_{Q,\rho} + \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \tilde{A}_{Q,\mu} \partial_\nu a_{Q,\rho} \quad (8)$$

where $\pm$ denotes the valley indices and $\tilde{A}_\mu = (\tilde{A}_0, \tilde{A}_x, \tilde{A}_y)$. Without any loss of generality, we assume $\tilde{A}_0 = 0$ for simplicity. The second term in Eq. (8) which is $j_Q^\mu \tilde{A}_{Q,\mu}$, is the minimal coupling between the induced gauge field and the current of quasiparticles. The Lagrangian of the whole system is then $\mathcal{L} = \mathcal{L}_K + \mathcal{L}_{K'}$. The corresponding action, $S = \int d^2x dt \mathcal{L}$, is known as mutual $U(1) \times U(1)$ Chern-Simons action. Although both $\mathcal{L}_K$ and $\mathcal{L}_{K'}$ break the TRS, their sum does not. This statement becomes obvious by considering the following transformations under the TRS:

$$\begin{aligned} K &\leftrightarrow K' \\ \tilde{A}_{K,\mu} &\leftrightarrow \tilde{A}_{K',\mu} \\ (a_{K,0}, \mathbf{a}_K) &\leftrightarrow (-a_{K',0}, \mathbf{a}_{K'}) \end{aligned} \quad (9)$$

The Lagrangian in Eq. (8) leads to the following equation of motion:

$$j_Q^\mu = \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \partial_\nu a_{Q,\rho} = \pm \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \partial_\nu \tilde{A}_{Q,\rho} \quad (10)$$

restoring $\hbar$ and e , the total current is

$$j^\mu = j_K^\mu + j_{K'}^\mu = \frac{2e^2}{\hbar} \epsilon^{\mu\nu\rho} \partial_\nu \tilde{A}_\rho \quad (11)$$

finally, the pseudo-magnetic and the pseudo-electric fields due to strain lead to the following induced currents

$$J_x = \frac{2e^2}{\hbar} \tilde{E}_{y,K}, \quad J_y = -\frac{2e^2}{\hbar} \tilde{E}_{x,K}, \quad n = \frac{2e^2}{\hbar} \tilde{B}_K \quad (12)$$

It is worth mentioning that the induced current creates a real vector potential, $\mathcal{A}_{EM}$ from Maxwell equation $\nabla \times \nabla \times \mathcal{A}_{EM} = \mu_0 \mathbf{J}$. This vector potential couples to the current of quasiparticles in the bulk $\mathbf{j}$, through $-e \int d^2x \mathcal{A}_{EM} \cdot \mathbf{j}$ term in the Lagrangian. Since both $\mathcal{A}_{EM}$ and $\mathbf{j}$ change sign under TRS, then the excess term is TRI and so is the total Lagrangian. It is worthwhile mentioning that $\mathcal{A}_{EM}$ is weak and since $C_K + C_{K'} = 0$, it does not generate edge current. Consequently, it can be neglected up to the first order of approximation.

One important consequence of these results is that the charge quantum Hall effect due to strains can serve as a new probe to detect topological insulator phase in graphene. In topological insulators, the charge current is zero, even in the presence of pseudo-electric field induced by strains. To see this, consider the Chern number of spin up electrons at valley K . We have $C_{K,\uparrow} = -C_{K,\downarrow} = \frac{1}{2}$. Therefore the current at K , is $J_K = J_{K,\uparrow} + J_{K,\downarrow} = \frac{e^2}{\hbar} (C_{K,\uparrow} + C_{K,\downarrow}) \tilde{E}_K = 0$. Similarly we have $J_{K'} = 0$ and therefore $J = J_K + J_{K'} = 0$. Consequently, we can measure the induced charge current in

a time reversal insulating graphene sample in response to strains. For topological insulators, on the other hand, this current will be zero, while for trivial insulators it is nonzero.




FIG. 2: (Color online) (Left): Proposed strain for a sample of graphene, $100\text{\AA} \times 100\text{\AA}$ with $P_B = 0.002\text{\AA}^{-1}$. The sample size is just used for demonstration. (Right): Direction of the charge current for different Dirac points in different edges.

We propose an experimental setup to measure CQHE based on recent developed techniques to grow graphene on a thin cuprates showing outstanding flexibility [26] subject to chemical doping using hydrogen-passivated [14]. The atomic displacements are given by $(u_x, u_y) = (-2P_B xy + P_E(t)y^2, P_B(x^2 + y^2))$ in the Cartesian coordinates (Fig. 2) where P_B and $P_E(t)$ are geometric parameters with unit of $1/\text{length}$ and $P_E(t)$ can be a time-varying periodic function, for instance $P_E(t) = P_B \cos(\omega t)$. The correspondence strain leads to a uniform magnetic field $\tilde{B}_z = 4\phi_0 c \beta P_B / a$ and pseudo-electric field $\tilde{E}_y = -2\phi_0 c \beta \omega \sin(\omega t) y / a$. The induced scalar potential, which is proportional to $u_{xx} + u_{yy}$, vanishes due the special introduced strain. Consider a strain geometry $P_B = 0.5 (\mu\text{m})^{-1}$ in a sample with size $0.4 \mu\text{m} \times 0.4 \mu\text{m}$, the pseudo-magnetic field yields $\tilde{B}_z \approx 9.0 \text{ T}$ which is large enough to observe the quantized quantum Hall Landau levels. Due to the finite width, there are finite edge modes in the zigzag boundary [27]. The zigzag boundary condition used here do not mix the K and K' valleys, leading to a clear distinction of the edge modes. Besides the edge modes, an external electric field can not induce pure current along the horizontal boundaries, however the pseudo-electric field constructed by $-\partial \tilde{A}(t)/\partial t$ creates easily a voltage along the boundaries demonstrated in Fig. 2. Importantly, using the parameters and $\omega = 60 \text{ MHz}$, the maximum pseudo-electric field reaches 50 V/m and then J_x would be about $4 \times 10^{-3} \text{ A/m}$. This induced current can be measured by experiments.

In summary, we have shown that CQHE can be seen even without the TRS breaking in a strained honeycomb structures which can be either artificial honeycomb lat-

tices or graphene nanoribbons. We have introduced staggered chemical potential to open a gap in the band structure together with a varying induced gauge field created by strain on the system. Furthermore we have shown that, even though that the whole system does not break the time reversal, the total current in the edge of the system is nonzero and we then observe CQHE. Finally, our theory can be investigated by an experimental setup that we have proposed.

We wish to thank A. Concha, M. Barkeshli and M. Vozmediano for useful discussions.

* Electronic address: vaezi@ipm.ir

† Electronic address: asgari@ipm.ir

- [1] K. V. Klitzing, G. Dorda, M. Pepper, Phys. Rev. Lett. **45**, 494 (1980); D. C. Tsui, H. L. Stormer, A. C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982); R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
- [2] K. S. Novoselov, *et al.* Nature **438** 197 (2005); Y. Zhang, *et al. ibid.* **438**, 201 (2005); X. Du *et al.*, *ibid.* **462**, 192 (2009); K.I. Bolotin *et al.*, *ibid.* **462**, 196 (2009).
- [3] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
- [4] A. Vaezi, arXiv:1105.0406; T. Neupert, *et al.*, arXiv:1012.4723; Y. -F. Wang, *et al.* arXiv: 1103.1686; E. Tang, J. -W. Mei, X. -G. Wen, arXiv: 1012.2930.
- [5] B. A. Bernevig, S.-C. Zhang, Phys. Rev. Lett. **96**, 106802 (2006); C. L. Kane, E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005).
- [6] D. Hsieh, D. Qian, L. Wray, Y. Xia, Y. S. Hor, R. J. Cava, M. Z. Hasan, Nature **452**, 970 (2008).
- [7] X. -G. Wen, Int. J. Mod. Phys. B **5**, 1641 (1991); X. -G. Wen, *Quantum Field Theory of Many-body Systems: From the Origin of Sound to an Origin of Light and Electrons*. Oxford University Press (2004).
- [8] M. A. H. Vozmediano, M. I. Katsnelson, F. Guinea, Phys. Reports **496**, 109 (2010).
- [9] N. Levy, *et al.*, Science **329**, 554 (2010).
- [10] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).
- [11] G. W. Semenoff, Phys. Rev. Lett. **53**, 2449 (1984).
- [12] F. Guinea, M. I. Katsnelson, and A. K. Geim, Nature Physics **6**, 30 (2010).
- [13] J. C. Slonczewski and P. R. Weiss, Phys. Rev. **109**, 272 (1958).
- [14] D. C. Elias *et al.*, Science **323**, 610 (2009).
- [15] M. Y. Han, B. Ozyilmaz, Y. Zhang, P. Kim, Phys. Rev. Lett. **98**, 206805 (2007).
- [16] M. Gibertini, *et al.*, Phys. Rev. B **79**, 241406 (2008).
- [17] Z. Y. Meng, *et al.*, Nature **464**, 847 (2010).
- [18] A. Vaezi, X. -G. Wen, arXiv:1010.5744.
- [19] A. Vaezi, X. -G. Wen, arXiv:1101.1662.
- [20] M.-T. Tran and K.-S. Kim, Phys. Rev. B **83**, 125416 (2011); J. Reuther, D. Abanin, R. Thomale, arXiv:1103.0859; S.-P. Kou *et al.*, arXiv:0910.2070.
- [21] It should be mentioned that in order to have a stable spin liquid we need to introduce the hopping to the next nearest neighbor as well.
- [22] Notice that these are not gauge fields, in the sense of gauge transformations i.e. assuming $\psi \rightarrow e^{i\alpha} \psi$, $\tilde{A}_\mu$ does

- not necessarily transform as $\tilde{A}_\mu \rightarrow \tilde{A}_\mu + \partial_\mu \alpha$.
- [23] H. Suzuura, T. Ando, Phys. Rev. B **65**, 235412 (2002);
J. L. Mañes, Phys. Rev. B **76** 045430 (2007) .
 - [24] F. Guinea, *et al.*, Phys. Rev. B **81**, 035408 (2010); N.
Abedpour, R. Asgari, F. Guinea, arXiv:1105.1026 .
 - [25] K.V. Zakharchenko, M.I. Katsnelson, and A. Fasolino,
Phys. Rev. Lett. **102**, 046808 (2009) .
 - [26] S. Bae, *et al.*, Nature nanotechnology **5**, 574 (2010); K.
S. Kim, *et al.*, Nature, **457**, 706 (2009) .
 - [27] T. Low, F. Guinea, Nano Lett. **10**, 3551 (2010) .